

The covariant Lyapunov tensor and the Lyapunov stability with respect to time-dependent Riemannian metrics

GENNADI SARDANASHVILY[†]

[†] Department of Theoretical Physics, Physics Faculty, Moscow State University, 117234 Moscow, Russia; E-mail: sard@grav.phys.msu.su

Abstract. We show that any solution of a smooth first order dynamic equation can be made Lyapunov stable at will by the choice of an appropriate time-dependent Riemannian metric.

1 Introduction

The Lyapunov matrix of a first order dynamic equation is defined as the coefficient matrix of the variation equation, and its properties are used as a criteria of stability of solutions of this equation [1, 4]. The problem is that the Lyapunov matrix is not a tensor under coordinate transformations, unless they are linear and time-independent. For instance, any first order dynamic equation defines a coordinate atlas such that its solution is constant on each coordinate chart (see Proposition 1 below) and, consequently, the Lyapunov matrix vanishes.

We introduce the covariant Lyapunov tensor of a first order dynamic equation and study the Lyapunov stability of solutions when this tensor is negative-definite. The covariant Lyapunov tensor essentially depends on the choice of a Riemannian metric. We show that, if a dynamic equation is defined by a complete vector field, the Lyapunov exponent of its solution can be made equal to any real number with respect to the appropriate time-dependent Riemannian metric. It follows that chaos in dynamical systems described by smooth (C^∞) first order dynamic equations can be characterized in full by time-dependent Riemannian metrics.

2 Geometry of first order dynamic equations in non-autonomous mechanics

Let $\mathbb{R}$ be the time axis provided with the Cartesian coordinate t and transition functions $t' = t + \text{const}$. In geometric terms, a (smooth) first order dynamic equation in non-autonomous mechanics is defined as a vector field γ on a smooth fibre bundle

$$\pi : Y \rightarrow \mathbb{R} \tag{1}$$

which obeys the condition $\gamma|dt = 1$ [5], i.e.,

$$\gamma = \partial_t + \gamma^\lambda \partial_\lambda. \quad (2)$$

The associated first order dynamic equation takes the form

$$\dot{t} = 1, \quad \dot{y}^\lambda = \gamma^\lambda(t, y^\mu) \partial_\lambda, \quad (3)$$

where $(t, y^\lambda, \dot{t}, \dot{y}^\lambda)$ are holonomic coordinates on TY . Its solutions are trajectories of the vector field γ (2). They assemble into a (regular) foliation $\mathcal{F}$ of Y . Equivalently, γ (2) is defined as a connection on the fibre bundle (1).

A fibre bundle Y (1) is trivial, but it admits different trivializations

$$Y \cong \mathbb{R} \times M, \quad (4)$$

distinguished by fibrations $Y \rightarrow M$. Moreover, Proposition 1 below appeals to an atlas of Y as a fibred manifold. If there is a trivialization (4) such that, with respect to the associated coordinates, the components γ^λ of the connection γ (2) are independent of t , one says that γ is a conservative first order dynamic equation on M .

Proposition 1. *Given a first order dynamic equation γ , there exists an atlas $\Psi = \{(U; t, y^a)\}$ of a fibred manifold $Y \rightarrow \mathbb{R}$ with time-independent transition functions $y'^a(y^b)$ such that any solution s of γ on each chart $(U; t, y^a)$ reads*

$$s^a(t) = \text{const.}, \quad t \in \pi(U) \subset \mathbb{R}.$$

Proof. The atlas Ψ is an atlas of adapted coordinates for the foliation $\mathcal{F}$ of trajectories of the vector field (2) [7]. $\square$

Proposition 2. *Let the vector field γ (2) be complete, i.e., there is a unique global solution of the dynamic equation γ through each point of Y . Then there is a trivialization (4) of Y such that any solution s of γ reads*

$$s^a(t) = \text{const.}, \quad t \in \mathbb{R},$$

with respect to associated bundle coordinates (t, y^a) .

Proof. If γ is complete, the foliation $\mathcal{F}$ of its trajectories is a fibration of Y along these trajectories onto any fibre of Y , e.g., $Y_{t=0} \cong M$. This fibration yields a desired trivialization [5, 6]. $\square$

One can think of the coordinates (t, y^a) in Proposition 2 as being the initial date coordinates because all points of the same trajectory differ from each other only in the temporal coordinate.

Let us consider the canonical lift $V\gamma$ of the vector field γ (2) onto the vertical tangent bundle VY of $Y \rightarrow \mathbb{R}$. With respect to the holonomic bundle coordinates $(t, y^\lambda, \bar{y}^\lambda)$ on VY , it reads

$$V\gamma = \gamma + \partial_\mu \gamma^\lambda \bar{y}^\mu \bar{\partial}_\lambda, \quad \bar{\partial}_\lambda = \frac{\partial}{\partial \bar{y}^\lambda}. \quad (5)$$

This vector field obeys the condition $V\gamma|dt = 1$, and defines the first order dynamic equation

$$\dot{t} = 1, \quad \dot{y}^\lambda = \gamma^\lambda(t, y^\nu), \quad (6a)$$

$$\dot{\bar{y}}^\lambda = \partial_\mu \gamma^\lambda(t, y^\nu) \bar{y}^\mu \quad (6b)$$

on VY . The equation (6a) coincides with the initial one (3). The equation (6b) is the well-known variation equation. Substituting a solution s of the initial dynamic equation (6a) into (6b), one obtains a linear dynamic equation whose solutions $\bar{s}$ are Jacobi fields of the solution s . In particular, if $Y \rightarrow \mathbb{R}$ is a vector bundle, there is the canonical splitting $VY \cong Y \times Y$ and the morphism $VY \rightarrow Y$ so that $s + \bar{s}$ obeys the initial dynamic equation (6a) modulo the terms of order > 1 in $\bar{s}$.

Remark that, if γ is the Hamilton equation for a Hamiltonian form H on

$$Y = V^*Q \rightarrow Q \rightarrow \mathbb{R}$$

in time-dependent Hamiltonian mechanics, the variation equation $V\gamma$ is also the Hamilton equation for the Hamiltonian form on V^*VQ which is the canonical tangent lift of H onto $VV^*Q = V^*VQ$ [2, 5, 6].

3 The covariant Lyapunov tensor

The collection of coefficients

$$l_\mu^\lambda = \partial_\mu \gamma^\lambda \quad (7)$$

of the variation equation (6b) is called the Lyapunov matrix. Clearly, it is not a tensor under bundle coordinate transformations of the fibre bundle Y (1). Let us bring l_μ^λ into a covariant tensor.

The stability conditions of Lyapunov for solutions of a first order dynamic equation involve the notion of a distance between different solutions at an instant t . Therefore, let a fibre bundle $Y \rightarrow \mathbb{R}$ be provided with a Riemannian fibre metric g , defined as a section of the symmetrized tensor product $\overset{2}{\vee} V^*Y \rightarrow Y$ of the vertical cotangent bundle

V^*Y of $Y \rightarrow \mathbb{R}$. With respect to the holonomic coordinates $(t, y^\lambda, \bar{y}_\lambda)$ on V^*Y , it takes the coordinate form

$$g = \frac{1}{2}g_{\alpha\beta}(t, y^\lambda)\bar{d}y^\alpha \vee \bar{d}y^\beta, \quad (8)$$

where $\{\bar{d}y^\mu\}$ are the holonomic fibre bases for V^*Y .

Given a first order differential equation γ , let

$$V^*\gamma = \gamma - \partial_\mu \gamma^\lambda \bar{y}_\lambda \bar{\partial}^\mu, \quad \bar{\partial}^\mu = \frac{\partial}{\partial \bar{y}_\mu}. \quad (9)$$

be the canonical lift of the vector field γ (2) onto V^*Y . It is a connection on $V^*Y \rightarrow \mathbb{R}$. Let us consider the Lie derivative $\mathbf{L}_\gamma g$ of the Riemannian fibre metric g along the vector field $V\gamma$ (9). It reads

$$L_{\alpha\beta} = (D_t g)_{\alpha\beta} = \partial_t g_{\alpha\beta} + \gamma^\lambda \partial_\lambda g_{\alpha\lambda} + \partial_\alpha \gamma^\lambda g_{\lambda\beta} + \partial_\beta \gamma^\lambda g_{\alpha\lambda}. \quad (10)$$

This is a tensor with respect to any bundle coordinate transformation of the fibre bundle (1). We agree to call it the covariant Lyapunov tensor. If g is an Euclidean metric, it comes to symmetrization

$$L_{\alpha\beta} = \partial_\alpha \gamma^\beta + \partial_\beta \gamma^\alpha = l_\alpha^\beta + l_\beta^\alpha$$

of the Lyapunov matrix (7).

Let us point the following two properties of the covariant Lyapunov tensor.

(i) Written with respect to the atlas Ψ in Proposition 1, the covariant Lyapunov tensor is

$$L_{ab} = \partial_t g_{ab}. \quad (11)$$

(ii) Given a solution s of the dynamic equation γ and a solution $\bar{s}$ of the variation equation (6b), we have

$$L_{\alpha\beta}(t, s^\lambda(t))\bar{s}^\alpha \bar{s}^\beta = \frac{d}{dt}(g_{\alpha\beta}(t, s^\lambda(t))\bar{s}^\alpha \bar{s}^\beta). \quad (12)$$

The definition of the covariant Lyapunov tensor (10) depends on the choice of a Riemannian fibre metric on the fibre bundle Y .

Proposition 3. *If the vector field γ is complete, there is a Riemannian fibre metric on Y such that the covariant Lyapunov tensor vanishes everywhere.*

Proof. Let us choose the atlas of the initial date coordinates in Proposition 2. Using the fibration $Y \rightarrow Y_{t=0}$, one can provide Y with a time-independent Riemannian fibre metric

$$g_{ab}(t, y^a) = h(t)g_{ab}^0(y^a) \quad (13)$$

where $g_{ab}^0(y^a)$ is a Riemannian metric on the fibre $Y_{t=0}$ and $h(t)$ is a positive smooth function on $\mathbb{R}$. The covariant Lyapunov tensor with respect to the metric (13) is

$$L_{ab} = \partial_t h g_{ab}.$$

Putting $h(t) = 1$, we obtain $L = 0$. □

4 The local Lyapunov stability

With the covariant Lyapunov tensor (10), we obtain the following variant of the stability condition of Lyapunov.

By the instantwise distance $\rho_t(s, s')$ between two solutions s and s' of the dynamic equation γ at an instant t is meant the distance between the points $s(t)$ and $s'(t)$ in the Riemannian space $(Y_t, g(t))$. Recall that a solution s of a first order dynamic equation g is said to be locally Lyapunov stable at an instant $t \in \mathbb{R}$ if, for any $\varepsilon > 0$, there is $\delta > 0$ such that $\rho_t(s, s') < \delta$ implies $\rho_{t'}(s, s') < \varepsilon$ for each t' which belongs to some half closed interval $[t, .)$. Being locally Lyapunov stable with respect to g , a solution is so with respect to any Riemannian fibre metric on Y .

Proposition 4. *Let the covariant Lyapunov tensor L (10) at a point $y \in Y$ be a negative-definite bilinear form on $V_y Y$. Then there is an open neighbourhood U_y of y and an open tubular neighbourhood $U_s \subset U_y$ of the trajectory s through y in U_y such that*

$$\rho_{t'>t}(s, s') < \rho_t(s, s') \tag{14}$$

for any $t' \in [t, .) \subset \pi(U_s)$ and s' crossing U_s .

Proof. Since the condition and the statement of Proposition 4 are coordinate-independent, let us choose a chart $(U; t, y^a)$ of the bundle atlas Ψ in Proposition 1 which cover the point y . With respect to these coordinates, the covariant Lyapunov tensor takes the form (11). Restricted to U , the leaf s of the foliation $\mathcal{F}$ is an imbedded submanifold, and has an open tubular neighbourhood W , i.e., any trajectory in U intersecting W does not leave W . There is an open neighbourhood $U_y \subset U$ of the point $y \in Y$ where the Lyapunov tensor L (10) holds negative-definite. Put $W' = W \cap U_y$. The foliation $\mathcal{F}$ restricted to the tubular W' defines its fibration

$$\zeta : W' \rightarrow W' \cap Y_{\pi(y)}$$

and the corresponding trivialization

$$W' \cong \pi(W') \times (W' \cap Y_{\pi(y)}). \tag{15}$$

There is an open neighbourhood $V \subset Y_{\pi(y)} \cap W'$ of y in $Y_{\pi(y)}$ which can be provided with the normal coordinates (x^a) defined by the Riemannian metric g in $Y_{\pi(y)}$ and centralized at y . Let us consider the open tubular $U_s = \zeta^{-1}(V)$. It is a subbundle of the trivial bundle (15) endowed with the coordinates (t, x^a) , $t \in (,)$. Without a loss of generality, put $t = \pi(y) = 0$. With respect to these coordinates, the solution s in U_s reads $s^a(t) = 0$. Let

$$s'^a(t) = u^a = \text{const.}, \quad t \in (,) = \pi(U_s), \quad (16)$$

be another solution crossing U_s . The instantwise distance $\rho_t(s, s')$ between solutions s and s' is the distance between the points $(t, 0)$ and (t, u) in the Riemannian space $(Y_t, g(t))$. This distance does not exceed the length

$$\bar{\rho}_t(s, s') = \left[\int_0^1 g_{ab}(t, \tau u^c) u^a u^b d\tau \right]^{1/2} \quad (17)$$

of the curve

$$x^a = \tau u^a, \quad \tau \in [0, 1] \quad (18)$$

in the Riemannian space $(Y_t, g(t))$. At the same time, we have

$$\rho_{t=0}(s, s') = \bar{\rho}_{t=0}(s, s')$$

The temporal derivative of the function $\bar{\rho}_t(s, s')$ (17) reads

$$\partial_t \bar{\rho}_t(s, s') = \frac{1}{2(\bar{\rho}_t(s, s'))^{1/2}} \int_0^1 \partial_t g_{ab}(t, \tau u^c) u^a u^b d\tau. \quad (19)$$

Since the bilinear form $\partial_t g_{ab} = L_{ab}$ is negative-definite at all points of the curve (18), the derivative (19) at all points $t \in (,)$ is also negative. Hence, we obtain

$$\rho_{t'>0}(s, s') < \bar{\rho}_{t'>0}(s, s') < \bar{\rho}_0(s, s') = \rho_0(s, s').$$

□

The inequality (14) shows that a solution of a first order differential equation which obeys the condition of Proposition 4 is locally Lyapunov stable at an instant t . Moreover, one can say that it is isometrically stable at t , and is so in some neighbourhood of t . Given another Riemannian fibre metric on Y , this solution remains locally Lyapunov stable, but need not be isometrically stable.

5 The asymptotic Lyapunov stability

Let γ (2) be a complete vector field. With a minor modification of the proof of Proposition 4, one can state the following.

Proposition 5. *Let s be a solution of the first order dynamic equation γ . If there exists an open tubular neighbourhood U_s of the trajectory s where the covariant Lyapunov tensor (10) is negative-definite at all instants t exceeding some t_0 , then*

$$\lim_{t' \rightarrow \infty} [\rho_{t'}(s, s') - \rho_t(s, s')] < 0 \quad (20)$$

for any $t > t_0$ and any solution $s' \subset U_s$.

One can say that a solution s in Proposition 5 obeys the isometric asymptotic stability of Lyapunov. Of course, it is also asymptotically stable, but so is any solution of a smooth dynamic equation as follows.

Proposition 6. *Any solution of any first order dynamic equation defined by a complete vector field (2) is Lyapunov stable at each instant of time and is asymptotically Lyapunov stable.*

Proof. These properties obviously hold with respect to the Riemannian fibre metric (13) in Proposition 3 where $h = 1$, and so do with respect to any Riemannian fibre metric on Y . $\square$

It follows that a solution of a first order dynamic equation is Lyapunov unstable only if this equation is given by a non-complete vector field γ (2). In this case, there exists a local (but not necessarily global) Riemannian fibre metric on Y such that a solution s of γ is locally stable at a given instant t .

One can improve Proposition 6 as follows.

Proposition 7. *Let λ be a real number. Given a dynamic equation γ defined by a complete vector field γ (2) and its solution s , there is a Riemannian fibre metric on Y such that the Lyapunov spectrum reduces to λ .*

Proof. Recall that the (upper) Lyapunov exponent of a solution s' with respect to a solution s is defined as the limit

$$K(s, s') = \lim_{t \rightarrow \infty} \frac{1}{t} \ln(\rho_t(s, s')). \quad (21)$$

Let provide the fibre bundle Y with the Riemannian fibre metric (13) in Proposition 3 where $h = \exp(\lambda t)$. A simple computation shows that the Lyapunov exponent (21) with respect to this metric is exactly λ . $\square$

If the upper limit

$$\lim_{\rho_{t=0}(s,s') \rightarrow 0} K(s, s') = \lambda$$

is negative, the solution s is said to be exponentially Lyapunov stable. If there exists at least one positive Lyapunov exponent, one speaks on chaos in a dynamical system [3]. Proposition 7 shows that chaos in smooth dynamical systems can be characterized in full by time-dependent Riemannian metrics.

References

- [1] Gallavotti G 1983 *The Elements of Mechanics* (Berlin: Springer-Verlag)
- [2] Giachetta G, Mangiarotti L and Sardanashvily G 1999 *J. Math. Phys.* **40** 1376
- [3] Gutzwiller M 1990 *Chaos in Classical and Quantum Mechanics* (Berlin: Springer)
- [4] Hirsch M and Smale S 1974 *Differential Equations, Dynamical Systems, and Linear Algebra* (New York: Academic Press)
- [5] Mangiarotti L and Sardanashvily G 1998 *Gauge Mechanics* (Singapore: World Scientific)
- [6] Mangiarotti L and Sardanashvily G 2000 *Connections in Classical and Quantum Field Theory* (Singapore: World Scientific)
- [7] Reinhart B 1983 *Differential Geometry and Foliations* (Berlin: Springer-Verlag, Berlin).